

International Union of Pure and Applied Chemistry
Division VIII Chemical Nomenclature and Structure Representation Division
***Nomenclature of Organic Chemistry. IUPAC Recommendations
and Preferred Names 2013.***

Prepared for publication by Henri A. Favre and Warren H. Powell,
Royal Society of Chemistry, ISBN 978-0-85404-182-4
School of Biological and Chemical Sciences, Queen Mary University of London,
Mile End Road, London, E1 4NS, UK

Prepared by Dr G P Moss <g.p.moss@qmul.ac.uk>

All noted corrections have been incorporated in this version. Otherwise the text is as close to the printed book as possible.

Contents

| Section | Title |
|---------|--|
| | MEMBERSHIP OF THE IUPAC COMMISSION ON NOMENCLATURE OF ORGANIC CHEMISTRY (CNOC) (1993- 2001) |
| | MEMBERSHIP OF THE IUPAC DIVISION OF CHEMICAL NOMENCLATURE AND STRUCTURE REPRESENTATION (2002-2013) |
| | PREFACE |
| | ACKNOWLEDGEMENTS |
| | CHANGES FROM THE 1979 EDITION, THE 1993 GUIDE, AND OFFICIAL PUBLICATIONS FROM 1993 THROUGH 2002 INCLUDED IN THE 2013 EDITION OF THE IUPAC NOMENCLATURE OF ORGANIC CHEMISTRY, 2013 EDITION |
| 1. | Scope of the recommendations in the 2013 edition |
| 2. | Skeletal replacement ('a') nomenclature |
| 3. | Substitutive nomenclature |
| 4. | Multiplicative nomenclature |
| 5. | Hydro or dehydro prefixes |
| 6. | Suffixes |
| 7. | Prefixes |
| 8. | Locants |
| 9. | Acyclic systems |
| 10. | Hantzsch-Widman system |
| 11. | Polycyclic ring systems |
| 12. | Seniority |
| 13. | Adducts |
| 14. | Enclosing marks |
| 15. | Esters |
| 16. | Amides |
| 17. | Inorganic parent structures |
| 18. | Radicals and ions |

19. Isotopically modified compounds

GLOSSARY

P-1 GENERAL PRINCIPLES, RULES, AND CONVENTIONS

- P-10 Introduction**
- P-11 Scope of nomenclature for organic compounds**
- P-12 Preferred, Preselected, and Retained IUPAC Names**
- P-12.1 Preferred IUPAC Names
- P-12.2 Preselected Names
- P-12.3 Retained Names
- P-12.4 Methodology
- P-13 Operations in Nomenclature of Organic Compounds**
- P-13.1 The Substitutive Operation
- P-13.2 The Replacement Operation
- P-13.3 The Additive Operation
- P-13.3.1 By an additive prefix
- P-13.3.2 By an additive suffix
- P-13.3.3 By a separate word
- P-13.4 The Subtractive Operation
- P-13.4.1 By a suffix
- P-13.4.2 By a change in ending
- P-13.4.3 By the prefixes 'dehydro' and 'nor'
- P-13.5 The Conjunctive Operation
- P-13.5.1 By placing a multiplicative prefix 'bi', 'ter', 'quater', etc.
- P-13.5.2 By juxtaposition of component names (conjunctive nomenclature)
- P-13.5.3 Ring formation
- P-13.6 The Multiplicative Operation
- P-13.7 The Fusion Operation
- P-13.8 Operations Used Only in the Nomenclature of Natural Products
- P-13.8.1 By the prefixes 'de' and 'des'
- P-13.8.2 By the prefix 'anhydro'
- P-14 General Rules**
- P-14.0 Introduction
- P-14.1 Bonding Number
- P-14.1.1 Definition
- P-14.1.2 Standard bonding numbers
- P-14.1.3 Nonstandard bonding numbers (λ -convention)
- P-14.2 Multiplicative Prefixes
- P-14.2.1 Basic multiplicative prefixes
- P-14.2.2 Numerical terms for compound or complex features
- P-14.2.3 Multiplicative prefixes for naming assemblies of identical units
- P-14.3 Locants

| | |
|-------------|--|
| P-14.3.1 | Types of locants |
| P-14.3.2 | Position of locants |
| P-14.3.3 | Citation of locants |
| P-14.3.4 | Omission of locants |
| P-14.3.5 | Lowest set of locants |
| P-14.4 | Numbering |
| P-14.5 | Alphanumerical Order |
| P-14.5.1 | Simple prefixes |
| P-14.5.2 | The name of a prefix for a substituent |
| P-14.5.3 | When an alphanumerical ordering is required |
| P-14.5.4 | When two or more prefixes consist of identical Roman letters |
| P-14.6 | Nonalphanumerical Order |
| P-14.7 | Indicated and 'Added Indicated' Hydrogen |
| P-14.7.1 | Indicated hydrogen |
| P-14.7.2 | 'Added indicated hydrogen' |
| P-14.8 | Adducts |
| P-14.8.1 | Organic adducts |
| P-14.8.2 | Mixed organic – inorganic adducts |
| P-15 | Types of Nomenclature |
| P-15.0 | Introduction |
| P-15.1 | Substitutive Nomenclature |
| P-15.1.1 | Parent hydrides |
| P-15.1.2 | Functional parent compounds |
| P-15.1.3 | Suffixes |
| P-15.1.4 | Position of the endings 'ane', 'ene', and 'yne' |
| P-15.1.5 | Prefixes and their order in names |
| P-15.1.6 | Other components of substitutive names |
| P-15.1.7 | Construction of substitutive names |
| P-15.1.8 | Substitution rules for parent structures with retained names |
| P-15.2 | Functional Class Nomenclature |
| P-15.2.0 | Introduction |
| P-15.2.1 | Functional class nomenclature using substituent group names |
| P-15.2.2 | Functional class nomenclature using functional modifiers |
| P-15.2.3 | Seniority of functional classes |
| P-15.2.4 | Polyfunctional compounds |
| P-15.3 | Multiplicative Nomenclature |
| P-15.3.0 | Introduction |
| P-15.3.1 | General methodology |
| P-15.3.2 | Construction of multiplicative names |
| P-15.3.3 | Identical structural units linked by unsymmetrical multiplicative groups |
| P-15.3.4 | Limitations of multiplicative nomenclature |
| P-15.4 | Skeletal Replacement ('a') Nomenclature |

| | |
|-------------|---|
| P-15.4.0 | Introduction |
| P-15.4.1 | General rules |
| P-15.4.2 | Skeletal replacement ('a') nomenclature |
| P-15.4.3 | Skeletal replacement ('a') nomenclature for acyclic parent hydrides |
| P-15.5 | Functional Replacement Nomenclature |
| P-15.5.1 | Definition |
| P-15.5.2 | General methodology |
| P-15.5.3 | Scope of functional replacement nomenclature |
| P-15.6 | Conjunctive Nomenclature |
| P-15.6.1 | Name formation |
| P-15.6.2 | Limitations of conjunctive nomenclature |
| P-15.6.3 | Analysis of conjunctive name construction |
| P-16 | Name Writing |
| P-16.0 | Introduction |
| P-16.1 | Spelling |
| P-16.2 | Punctuation |
| P-16.2.1 | Commas |
| P-16.2.2 | Full stops |
| P-16.2.3 | Colons and semicolons |
| P-16.2.4 | Hyphens |
| P-16.2.5 | Spaces |
| P-16.3 | Multiplicative prefixes 'di', 'tri', etc. vs. 'bis', 'tris', etc. |
| P-16.3.1 | Multiplicative prefixes |
| P-16.3.2 | General methodology |
| P-16.3.3 | The basic numerical prefixes 'di', 'tri', 'tetra', etc. |
| P-16.3.4 | Parentheses (round brackets) |
| P-16.3.5 | The numerical prefixes 'bis', 'tris', 'tetrakis', etc. |
| P-16.3.6 | The prefixes 'bis', 'tris', 'tetrakis', etc. |
| P-16.4 | Other Numerical Terms |
| P-16.4.1 | The numerical prefixes 'bi', 'ter', 'quater', etc. |
| P-16.4.2 | The prefix 'mono' |
| P-16.4.3 | The term 'sesqui' |
| P-16.5 | Enclosing Marks |
| P-16.5.1 | Parentheses (also called curves or round brackets) |
| P-16.5.2 | Brackets (also called square brackets) |
| P-16.5.3 | Braces (also called curly brackets) |
| P-16.5.4 | Multiple types of enclosing marks |
| P-16.6 | Italicization |
| P-16.6.1 | Lower case italic letters |
| P-16.6.2 | Italicized elements symbols, such as O, N, As |
| P-16.6.3 | The italic element symbol <i>H</i> |
| P-16.6.4 | Italic terms, syllables, and capital Roman letters |

| | |
|----------|--|
| P-16.7 | Elision of Vowels |
| P-16.7.1 | Vowels systematically elided |
| P-16.7.2 | There is no elision of terminal vowels |
| P-16.8 | Addition of Vowels |
| P-16.8.1 | For euphonic reasons (the vowel ‘o’) |
| P-16.8.2 | For euphonic reasons (the letter ‘a’) |
| P-16.9 | Primes |
| P-16.9.1 | Primes (′), double primes (″), triple primes (″″), etc., with nitrogen |
| P-16.9.2 | Superscript arabic numbers, |
| P-16.9.3 | Primes (′), double primes (″), triple primes (″″), etc., with superscripted numerical locants |
| P-16.9.4 | Primes (′), double primes (″), triple primes (″″), etc., used to differentiate identical locants |
| P-16.9.5 | Other uses of primes |
| P-16.9.6 | Multiple primes |

P-2 PARENT HYDRIDES

| | |
|-------------|--|
| P-20 | Introduction |
| P-21 | Mononuclear and Acyclic Polynuclear Parent Hydrides |
| P-21.1 | Mononuclear Parent Hydrides |
| P-21.1.1 | Mononuclear parent hydrides with standard bonding numbers |
| P-21.1.2 | Mononuclear parent hydrides with nonstandard bonding numbers |
| P-21.2 | Acyclic Polynuclear Parent Hydrides |
| P-21.2.1 | Hydrocarbons |
| P-21.2.2 | Homogeneous acyclic parent hydrides other than hydrocarbons and boron hydrides |
| P-21.2.3 | Heterogenous acyclic parent hydrides |
| P-21.2.4 | Acyclic parent hydrides containing heteroatoms with nonstandard bonding numbers |
| P-22 | Monocyclic Parent Hydrides |
| P-22.1 | Monocyclic Hydrocarbons |
| P-22.1.1 | Saturated monocyclic hydrocarbons |
| P-22.1.2 | Unsubstituted monocyclic hydrocarbon polyenes |
| P-22.1.3 | Parent hydrocarbons having retained names |
| P-22.2 | Heteromonocyclic Parent Hydrides |
| P-22.2.1 | Retained names of heteromonocycles |
| P-22.2.2 | Heteromonocyclic parent hydrides with 3-10 membered rings (Hantzsch-Widman names) |
| P-22.2.3 | Heteromonocyclic parent hydrides named by skeletal replacement (‘a’) nomenclature |
| P-22.2.4 | Heteromonocycles with eleven or more members with the maximum number of noncumulative double bonds |
| P-22.2.5 | Homogeneous monocyclic parent hydrides other than boron hydrides |
| P-22.2.6 | Heteromonocyclic parent hydrides composed of repeating heterounits |

| | |
|-------------|--|
| P-22.2.7 | Heteromonocyclic parent hydrides having heteroatoms with nonstandard bonding numbers |
| P-23 | Polyalicyclic Parent Hydrides (Extended von Baeyer System) |
| P-23.0 | Introduction |
| P-23.1 | Definitions and Terminology |
| P-23.2 | Naming and Numbering of von Baeyer Hydrocarbons |
| P-23.2.1 | Selection of the main ring |
| P-23.2.2 | Naming bicyclic alicyclic hydrocarbons |
| P-23.2.3 | Numbering bicyclic alicyclic hydrocarbons |
| P-23.2.4 | Selection of the main bridge |
| P-23.2.5 | Naming and numbering tricyclic alicyclic hydrocarbons |
| P-23.2.6 | Naming and numbering polyalicyclic hydrocarbons |
| P-23.2.6.1 | Naming polycyclic alicyclic hydrocarbons |
| P-23.2.6.2 | Selection of the main bridge and secondary bridges |
| P-23.2.6.3 | Numbering of secondary bridges |
| P-23.2.6.4 | If there is a further choice for numbering the secondary bridges |
| P-23.3 | Heterogeneous Heterocyclic von Baeyer Parent Hydrides |
| P-23.3.1 | Heterogeneous heterocyclic von Baeyer systems |
| P-23.3.2 | When there is a choice for numbering |
| P-23.4 | Homogeneous Heterocyclic von Baeyer Parent Hydrides |
| P-23.5 | Heterogeneous Heterocyclic von Baeyer Parent Hydrides Composed of Alternating Heteroatoms |
| P-23.5.1 | Heterogeneous von Baeyer systems composed of alternating skeletal heteroatoms |
| P-23.5.2 | The prefixes 'Si-' or 'N-' preceded by the italic number '1' |
| P-23.5.3 | Silasesquioxanes, silasesquithianes, etc. |
| P-23.6 | Heterocyclic Polyalicyclic Parent Hydrides having Heteroatoms with Nonstandard Bonding Numbers |
| P-23.6.1 | The λ -convention |
| P-23.6.2 | When there is a choice for numbering |
| P-23.7 | Retained Names for von Baeyer Parent Hydrides |
| P-24 | Spiro Ring Systems |
| P-24.0 | Introduction |
| P-24.1 | Definitions |
| P-24.2 | Spiro Ring Systems with only Monocyclic Ring Components |
| P-24.2.0 | Introduction |
| P-24.2.1 | Monospiro alicyclic ring systems |
| P-24.2.2 | Linear polyspiro alicyclic ring systems |
| P-24.2.3 | Branched polyspiro alicyclic ring systems |
| P-24.2.4 | Heterocyclic spiro ring systems |
| P-24.3 | Monospiro Ring Systems Containing Two Identical Polycyclic Ring Components |
| P-24.3.1 | Monospiro ring systems consisting of two identical polycyclic ring |

| | |
|-------------|---|
| | components |
| P-24.3.2 | Where appropriate the maximum number of noncumulative double bonds |
| P-24.3.3 | If there is a choice for assigning primed locants |
| P-24.3.4 | Heterocyclic 'spirobi' ring systems named by skeletal replacement ('a') nomenclature |
| P-24.4 | Dispiro Ring Systems with Three Identical Polycyclic Ring Components Spiro-fused Together |
| P-24.4.1 | Dispiro ring systems with three identical polycyclic ring components |
| P-24.4.2 | If there is a choice for locants |
| P-24.4.3 | Three identical heterocyclic ring components spiro-fused together |
| P-24.5 | Monospiro Ring Systems with Different Components, at least One of which is a Polycyclic Ring System |
| P-24.5.1 | Monospiro ring systems with different ring components |
| P-24.5.2 | Monospiro ring systems with different ring components at least one being a polycyclic ring and at least one ring component requiring the use of skeletal replacement ('a') nomenclature |
| P-24.5.3 | Alphanumerical order |
| P-24.5.4 | Ring systems modified by skeletal replacement ('a') nomenclature |
| P-24.6 | Unbranched Polyspiro Ring Systems with Different Components, One being a Polycyclic Ring System |
| P-24.6.1 | If there is a choice of locants |
| P-24.7 | Branched Polyspiro Ring Systems |
| P-24.7.1 | A central ring component is spiro-fused to three or more identical terminal ring components |
| P-24.7.2 | Two or more different terminal ring components spiro-fused to a central component |
| P-24.7.3 | If no decision can be attained by application of P-24.7.2 |
| P-24.7.4 | Additional components are spiro-fused to a branched polyspiro compound |
| P-24.8 | Spiro Ring Systems Containing Atoms with Nonstandard Bonding Numbers |
| P-24.8.1 | Spiro ring systems with only monocyclic ring components |
| P-24.8.2 | Monospiro ring systems with two identical polycyclic ring components |
| P-24.8.3 | Spiro ring systems with three identical components and one nonstandard spiro atom |
| P-24.8.4 | Monospiro ring systems with different ring components, at least one of which is a polycyclic ring with a nonstandard spiro atom |
| P-24.8.5 | Unbranched polyspiro ring systems with different ring components, at least one of which is a polycyclic ring with at least one nonstandard spiro atom |
| P-24.8.6 | Branched spiro ring systems with at least one polycyclic ring component |
| P-25 | Fused and Bridged Fused Ring Systems |
| P-25.0 | Introduction |
| P-25.1 | Names of Hydrocarbon Parent Ring Components |
| P-25.1.1 | Retained names for hydrocarbons |

| | |
|--------------|---|
| P-25.1.2 | Systematically named hydrocarbon parent components |
| P-25.1.2.1 | Polyacenes |
| P-25.1.2.2 | Polyaphenes |
| P-25.1.2.3 | Polyalenes |
| P-25.1.2.4 | Polyphenylenes |
| P-25.1.2.5 | Polynaphthylenes |
| P-25.1.2.6 | Polyhelicenes |
| P-25.1.2.7 | Ace...ylenes |
| P-25.2 | Names of Heterocyclic Parent Ring Components |
| P-25.2.1 | Retained names for heterocycles |
| P-25.2.2 | Systematically named heterocyclic components |
| P-25.2.2.1 | Heteromonocyclic parent components |
| P-25.2.2.2 | Heteranthrene components |
| P-25.2.2.3 | Pheno...ine components |
| P-25.2.2.4 | Heteromonocyclic components fused to a benzene ring |
| P-25.3 | Constructing Fusion Names |
| P-25.3.1 | Definitions, terminology, and general principles |
| P-25.3.2 | Constructing two-component fusion names |
| P-25.3.3 | Numbering of fused ring systems |
| P-25.3.4 | Constructing polycomponent fusion names |
| P-25.3.4.1 | Three types of fusion names |
| P-25.3.4.2 | Constructing polycomponent fusion names |
| P-25.3.4.2.1 | Order of seniority for selecting parent components |
| P-25.3.4.2.2 | Order of seniority for selecting attached components |
| P-25.3.4.2.3 | Order of citation of fusion prefixes |
| P-25.3.4.2.4 | Order of seniority of locants (letters and numbers) |
| P-25.3.5 | Heteromonocyclics fused to a benzene ring |
| P-25.3.6 | Identical attached components |
| P-25.3.7 | Multiparent ring systems |
| P-25.3.8 | Omission of locants in fusion descriptors |
| P-25.4 | Bridged Fused Ring Systems |
| P-25.4.1 | Definitions and terminology |
| P-25.4.2 | Names of bridges |
| P-25.4.2.1 | Bivalent bridges |
| P-25.4.2.2 | Polyvalent bridges |
| P-25.4.2.3 | Composite bridges |
| P-25.4.3 | Naming bridged fused systems |
| P-25.4.4 | Numbering bridge atoms |
| P-25.4.5 | Order for numbering of bridges |
| P-25.5 | Limitations of Fusion Nomenclature: Three Components <i>ortho</i> and <i>peri</i> -fused Together |
| P-25.5.1 | Skeletal replacement ('a') nomenclature |

| | |
|-------------|--|
| P-25.5.2 | Selection of a less senior hydrocarbon parent component |
| P-25.5.3 | Use of bridging nomenclature |
| P-25.6 | Fused Ring Systems with Skeletal Atoms with Nonstandard Bonding Numbers |
| P-25.7 | Double Bonds, Indicated Hydrogen, and the δ -Convention |
| P-25.7.1 | Indicated hydrogen |
| P-25.7.2 | The δ -convention |
| P-25.8 | Parent Components in Decreasing Order of Seniority (Partial Lists) |
| P-25.8.1 | Partial list of heterocyclic parent components in decreasing order of seniority |
| P-25.8.2 | Partial list of hydrocarbon parent components in decreasing order of seniority |
| P-26 | Phane Nomenclature |
| P-26.0 | Introduction |
| P-26.1 | Concepts and Terminology |
| P-26.1.1 | Simplification and amplification |
| P-26.1.2 | Simplified skeleton of the phane parent hydride, simplified phane parent graph, simplified skeletal name, and skeletal locants |
| P-26.1.3 | Superatom and superatom locants |
| P-26.1.4 | Amplificant, amplification prefix, and amplificant locants |
| P-26.1.5 | Attachment atoms and attachment locants |
| P-26.1.6 | Phane parent skeleton, phane parent name, and phane parent hydrides |
| P-26.2 | Components of Phane Parent Names |
| P-26.2.1 | Simplified skeletal names |
| P-26.2.2 | Amplification prefixes |
| P-26.2.3 | Multiple identical amplificants |
| P-26.3 | Superatom Locants and Amplificant Attachment Locants |
| P-26.3.1 | Superatom locants |
| P-26.3.2 | Attachment locants |
| P-26.4 | Numbering of Phane Parent Hydride |
| P-26.4.1 | Numbering of phane parent skeletons and amplificants |
| P-26.4.2 | Numbering of simplified phane skeletons with respect to amplificant numbering |
| P-26.4.3 | Numbering of phane parent hydrides |
| P-26.5 | Skeletal Replacement ('a') Nomenclature in Phane Nomenclature |
| P-26.5.1 | Skeletal replacement ('a') nomenclature in simplified phane names |
| P-26.5.2 | Skeletal replacement ('a') nomenclature in amplificants |
| P-26.5.3 | Simultaneous skeletal replacement ('a') in simplified skeletal names and amplificants |
| P-26.5.4 | Numbering of heterophane parent hydrides with respect to heteroatoms |
| P-26.6 | Other Aspects of Phane Nomenclature |
| P-27 | Fullerenes |
| P-27.0 | Introduction |

| | |
|-------------|--|
| P-27.1 | Definitions |
| P-27.1.1 | Fullerenes |
| P-27.1.2 | Fulleranes |
| P-27.1.3 | Fulleroids |
| P-27.2 | Fullerene Names |
| P-27.2.1 | Systematic names |
| P-27.2.2 | Trivial names |
| P-27.2.3 | Preferred IUPAC names |
| P-27.3 | Numbering of Fullerenes |
| P-27.4 | Structurally Modified Fullerenes |
| P-27.4.0 | Introduction |
| P-27.4.1 | Homofullerenes |
| P-27.4.2 | Norfullerenes |
| P-27.4.3 | Secofullerenes |
| P-27.4.4 | Cyclofullerenes |
| P-27.4.5 | Combination of structure-modifying operations |
| P-27.5 | Replacement of Skeletal Atoms |
| P-27.5.1 | Fullerenes in which carbon atoms have been replaced by one or more heteroatoms |
| P-27.5.2 | When 'homo', 'nor', 'seco', or 'cyclo' prefixes co-occur with skeletal replacement terms, such as 'oxa' or 'aza' |
| P-27.6 | Addition of Rings and Ring Systems to Fullerenes |
| P-27.6.1 | Fullerenes ortho fused to organic rings or ring systems |
| P-27.6.2 | Bridged fullerenes |
| P-27.6.3 | Spiro fullerenes |
| P-27.7 | Other Aspects of Fullerene Nomenclature |
| P-28 | Ring Assemblies |
| P-28.0 | Introduction |
| P-28.1 | Definitions |
| P-28.2 | Ring Assemblies of Two Identical Cyclic Systems |
| P-28.2.1 | Ring assemblies with a single bond junction |
| P-28.2.2 | Ring assemblies with a double bond junction |
| P-28.2.3 | Indicated hydrogen of components in two-component ring assemblies |
| P-28.3 | Unbranched Ring Assemblies of Three through Six Identical Cyclic Systems |
| P-28.3.1 | Unbranched ring assemblies consisting of three or more identical cyclic systems |
| P-28.3.2 | Exceptions to the use of parent hydride names, ring assemblies of three through six benzene rings |
| P-28.4 | Ring Assemblies Composed of Identical Cyclic Systems Modified by Skeletal Replacement ('a') Nomenclature |
| P-28.4.1 | Assemblies composed of identical heterocyclic compounds |
| P-28.4.2 | Since hydrocarbon ring assemblies are the parent structures for the application of skeletal replacement ('a') nomenclature |

| | |
|-------------|--|
| P-28.5 | Ring Assembles Composed of More than Six Identical Cyclic Systems |
| P-28.6 | Branched Ring Assemblies of Identical Cyclic Systems |
| P-28.7 | Ring Assemblies of Nonidentical Cyclic Systems |
| P-29 | Prefixes Denoting Substituent Groups Derived from Parent Hydrides |
| P-29.0 | Introduction |
| P-29.1 | Definitions |
| P-29.1.1 | A simple substituent group |
| P-29.1.2 | A compound substituent group |
| P-29.1.3 | A complex substituent group |
| P-29.2 | General Methodology for Naming Substituent Groups Prefixes |
| P-29.3 | Systematic Prefixes for Simple Substituents Groups Derived from Saturated Parent Hydrides |
| P-29.3.1 | Substituent prefixes derived from mononuclear parent hydrides |
| P-29.3.2 | Substituent prefixes derived from acyclic parent hydrides |
| P-29.3.3 | Substituent prefixes derived from saturated cyclic parent hydrides |
| P-29.3.4 | Substituent prefixes derived from mancude parent hydrides |
| P-29.3.5 | Substituent prefixes derived from ring assemblies |
| P-29.3.6 | Substituent prefixes derived from phane systems |
| P-29.4 | Compound Substituent Groups |
| P-29.4.1 | Compound substituted substituent groups |
| P-29.4.2 | Compound catenated substituent groups |
| P-29.5 | Complex Substituent Groups |
| P-29.5.1 | Complex substituted substituent groups |
| P-29.5.2 | Concatenated complex substituent groups |
| P-29.6 | Retained Names for Prefixes of Simple Substituent Groups Derived from the Parent Hydrides Described in Chapter P-2 |
| P-29.6.1 | Retained prefixes that are preferred prefixes |
| P-29.6.2 | Retained prefixes that are not used as preferred prefixes |
| P-29.6.3 | Retained prefixes no longer recommended as approved prefixes |

P-3 CHARACTERISTIC (FUNCTIONAL) AND SUBSTITUENT GROUPS

| | |
|-------------|---|
| P-30 | Introduction |
| P-31 | Modification of the Degree of Hydrogenation of Parent Hydrides |
| P-31.0 | Introduction |
| P-31.1 | Substituent Groups Derived from Parent Hydrides with 'ene' or 'yne' endings |
| P-31.1.1 | General methodology |
| P-31.1.2 | Acyclic parent hydrides |
| P-31.1.3 | Monocyclic parent hydrides |
| P-31.1.4 | Bi- and polycyclic von Baeyer parent hydrides |
| P-31.1.5 | Spiro compounds |
| P-31.1.6 | Phane parent hydrides |
| P-31.1.7 | Ring assemblies of unsaturated components |

| | |
|--------------|---|
| P-31.2 | Substituent Groups Modified by the Prefixes 'hydro' or 'dehydro' |
| P-31.2.1 | The prefixes 'hydro/dehydro' |
| P-31.2.2 | General methodology |
| P-31.2.3 | The prefix 'hydro' |
| P-31.2.3.1 | Modification of the degree of hydrogenation of monocyclic mancude compounds |
| P-31.2.3.2 | Names of saturated heteromonocyclic compounds |
| P-31.2.3.3 | Saturation of double bonds in polycyclic mancude compounds |
| P-31.2.3.3.1 | Retained names of partially saturated polycyclic mancude compounds |
| P-31.2.3.3.2 | Polycyclic mancude compounds |
| P-31.2.3.3.3 | Spiro compounds |
| P-31.2.3.3.4 | Phane compounds |
| P-31.2.3.3.5 | Ring assemblies |
| P-31.2.4 | The prefix 'dehydro' |
| P-32 | Prefixes for Substituent Groups Derived from Parent Hydrides with a Modified Degree of Hydrogenation |
| P-32.0 | Introduction |
| P-32.1 | Substituent Groups Derived from Parent Hydrides with 'ene' or 'yne' endings |
| P-32.1.1 | Substituents derived from unsaturated acyclic compounds |
| P-32.1.2 | Monocyclic substituent groups |
| P-32.1.3 | Substituent groups derived from parent hydrides having a fixed numbering |
| P-32.2 | Substituent Groups Modified by the Prefix 'hydro' |
| P-32.2.1 | For heteromonocyclic parent hydrides |
| P-32.2.2 | For polycyclic mancude compounds |
| P-32.2.3 | With added indicated hydrogen |
| P-32.3 | Retained Names for Substituent Groups Derived from Unsaturated Acyclic Parent Hydrides |
| P-32.4 | Retained Names for Substituent Groups Derived from Partially Saturated Polycyclic Parent Hydrides |
| P-33 | Suffixes |
| P-33.0 | Introduction |
| P-33.1 | Definitions |
| P-33.2 | Functional Suffixes |
| P-33.2.1 | Basic functional suffixes |
| P-33.2.2 | Derived preferred and preselected suffixes |
| P-33.3 | Cumulative Suffixes |
| P-34 | Functional Parent Compounds |
| P-34.0 | Introduction |
| P-34.1 | Retained Functional Parent Compounds |
| P-34.1.1 | Organic functional parent compounds (arranged by characteristic group class) |

| | |
|-------------|---|
| P-34.1.2 | Organic functional parent compounds (arranged alphabetically) |
| P-34.1.3 | Organic functional parent compounds for general and specialized nomenclature |
| P-34.1.4 | Inorganic functional parent compounds |
| P-34.2 | Substituent Groups Related to Functional Parent Compounds |
| P-34.2.1 | Organic substituent groups (arranged by classes) |
| P-34.2.2 | Organic substituent groups (arranged alphabetically) |
| P-34.2.3 | Substituent group names derived from organic compounds used in general and specialized nomenclature |
| P-34.2.4 | Preselected substituent group names |
| P-35 | Prefixes Corresponding to Characteristic Groups |
| P-35.0 | Introduction |
| P-35.1 | General Methodology |
| P-35.2 | Simple Prefixes Denoting Characteristic Groups |
| P-35.2.1 | Retained traditional prefixes |
| P-35.2.2 | Substituents formed by subtracting one or more hydrogen atoms from mono and dinuclear parent hydrides |
| P-35.2.3 | Simple prefixes derived from functional parent compounds |
| P-35.3 | Compound Substituent Prefixes |
| P-35.3.1 | Names of compound prefixes derived from suffixes or functional parent compounds by substitution |
| P-35.3.2 | Names of compound prefixes derived from suffixes or functional parent compounds by concatenation |
| P-35.4 | Complex Substituent Prefixes |
| P-35.4.1 | Complex substituent prefixes by substitution |
| P-35.4.2 | Complex substituent prefixes by concatenation |
| P-35.5 | Mixed Substituent Prefixes |
| P-35.5.1 | Mixed substituent prefixes names using substitutive and additive operations |
| | P-4 RULES FOR NAME CONSTRUCTION |
| P-40 | Introduction |
| P-41 | Seniority Order for Classes |
| P-42 | Seniority Order for Acids |
| P-42.1 | Class 7a. Acids Expressed by Suffixes (Excludes Carbonic and Polycarbonic acids) |
| P-42.2 | Class 7b. Carbon Acids with no Substitutable Hydrogen Atoms |
| P-42.3 | Class 7c. Noncarbon Acids having Substitutable Hydrogen Atoms on the Central Atom |
| P-42.4 | Class 7d. Noncarbon Acids used to Generate Derivatives having Substitutable Hydrogen Atoms |
| P-42.5 | Class 7e. Other Monobasic 'oxo' Acids used as Functional Parents |
| P-43 | Seniority Order for Suffixes |
| P-43.0 | Introduction |

| | |
|--------------|--|
| P-43.1 | General Methodology of Functional Replacement |
| P-44 | Seniority order for parent structures |
| P-44.0 | Introduction |
| P-44.1 | Seniority Order for Parent Structures |
| P-44.1.1 | Maximum number of substituents corresponding to the principal characteristic group |
| P-44.1.2 | Senior atom in accordance with the seniority of classes |
| P-44.1.3 | Seniority order only for rings and ring systems |
| P-44.1.4 | Seniority among acyclic chains (the principal chain) |
| P-44.1.5 | Other criteria |
| P-44.2 | Seniority Order only for Rings and Ring Systems |
| P-44.2.1 | General criteria (other thanphanes) |
| P-44.2.1.1 | General methodology |
| P-44.2.1.2 | A heterocycle |
| P-44.2.1.3 | At least one nitrogen ring atom |
| P-44.2.1.4 | At least one heteroatom (in the absence of nitrogen) that occurs earlier from a list |
| P-44.2.1.5 | The greater number of rings |
| P-44.2.1.6 | Greater number of skeletal atoms |
| P-44.2.1.7 | Greater number of heteroatoms of any kind |
| P-44.2.1.8 | Greater number of heteroatoms that occur earlier from a list |
| P-44.2.2 | Criteria specific to a particular kind of ring or ring system |
| P-44.2.2.1 | Monocycles |
| P-44.2.2.2 | Polycyclic systems |
| P-44.2.2.2.1 | Spiro ring systems |
| P-44.2.2.2.2 | Cyclic phane systems |
| P-44.2.2.2.3 | Fused ring systems |
| P-44.2.2.2.4 | Bridged fused ring systems |
| P-44.2.2.2.5 | Von Baeyer ring systems |
| P-44.2.2.2.6 | Linear (acyclic)phanes |
| P-44.2.2.2.7 | Ring assemblies |
| P-44.3 | Seniority of Acyclic Chains (the Principal Chain) |
| P-44.3.1 | Greater number of acyclic heteroatoms of any kind |
| P-44.3.2 | Greater number of skeletal atoms |
| P-44.3.3 | Greater number of the most senior acyclic heteroatom |
| P-44.4 | Seniority Criteria Applicable to Rings, Ring Systems, or Acyclic Chains |
| P-44.4.1 | If P-44.1 through P-44.3, does not effect a choice of senior parent structure |
| P-44.4.1.1 | Greater number of multiple bonds |
| P-44.4.1.2 | Greater number of double bonds |
| P-44.4.1.3 | One or more atoms with nonstandard bonding numbers |
| P-44.4.1.4 | Lowest locants for indicated hydrogen |
| P-44.4.1.5 | Lower locants for heteroatoms introduced by skeletal replacement |

| | |
|-------------|---|
| | (‘a’) nomenclature as a set |
| P-44.4.1.6 | Lower locants for heteroatoms introduced by skeletal replacement (‘a’) nomenclature in the order |
| P-44.4.1.7 | Lower locant(s) for carbon atoms at fusion sites |
| P-44.4.1.8 | Lower locants for characteristic groups cited as a suffix |
| P-44.4.1.9 | Lowest locants for points of attachment (if a substituent group) |
| P-44.4.1.10 | Lowest locants for endings or prefixes that express changes in the degree of hydrogenation, i.e., for ‘ene’ and ‘yne’ endings and ‘hydro/dehydro’ prefixes |
| P-44.4.1.11 | One or more isotopically modified atoms |
| P-44.4.1.12 | One or more stereogenic centers |
| P-45 | Selection of the Preferred IUPAC Name |
| P-45.0 | Introduction |
| P-45.1 | Multiplication of Identical Senior Parent Structures |
| P-45.1.1 | Multiplicative nomenclature is senior to substitutive nomenclature |
| P-45.1.2 | Choice between two or more parent structures |
| P-45.2 | Criteria Related to Number and Location of Substituent Groups |
| P-45.2.1 | Maximum number of substituent groups cited as prefixes |
| P-45.2.2 | Lower locant set for substituent groups cited as prefixes |
| P-45.2.3 | Lower locant set for substituent groups in order of citation in the name |
| P-45.3 | Criteria Related only to Substituents with Nonstandard Bonding Numbers, other Criteria being Equal |
| P-45.3.1 | Maximum number of substituent group(s) with the higher bonding number cited as prefixes |
| P-45.3.2 | Lower locant set for substituent group(s) with the higher bonding number cited as prefixes |
| P-45.4 | Criteria Related only to Isotopic Modification of Substituents |
| P-45.4.1 | Senior parent structure that has the lowest locant(s) for isotopically modified substituent groups |
| P-45.4.2 | Senior parent structure that has the lowest locant(s) for nuclides of higher atomic number |
| P-45.4.3 | Senior parent structure that has the lowest locant(s) for nuclides of higher mass number |
| P-45.5 | Criteria Related to Alphanumerical Order of Names |
| P-45.5.1 | Earlier in alphanumerical order |
| P-45.5.2 | When names based on alphanumerical order are the same |
| P-45.6 | Criteria Related only to Configuration |
| P-45.6.1 | Introduction |
| P-45.6.2 | General principles |
| P-45.6.3 | Alphabetic order of the stereochemical descriptors ‘R’ and ‘S’ |
| P-45.6.4 | Application of CIP sequence rules 4 and 5 |
| P-46 | The Principal Chain in Substituent Groups |
| P-46.0 | Introduction |
| P-46.1 | The Principal Substituent Chain |

| | |
|-----------|---|
| P-46.1.1 | Greater number of heteroatoms |
| P-46.1.2 | Greater number of skeletal atoms |
| P-46.1.3 | Greater number of heteroatoms in a set order |
| P-46.1.4 | Greater number of multiple bonds, then double bonds |
| P-46.1.5 | One or more atoms with nonstandard bonding numbers |
| P-46.1.6 | Lowest locants for heteroatoms |
| P-46.1.7 | Lowest locants for heteroatoms appearing first a set order |
| P-46.1.8 | Lowest locants for free valences of any kind in the order 'yl' > 'ylidene' > 'ylidyne' |
| P-46.1.9 | Lowest locant(s) for multiple bonds, then double bonds |
| P-46.1.10 | Lowest locant(s) for (an) atom(s) with nonstandard bonding number(s) |
| P-46.1.11 | Greatest number of substituents of any kind |
| P-46.1.12 | Lowest locants for substituents |
| P-46.1.13 | Lowest locants for the substituent(s) cited earlier in alphanumerical order |
| P-46.2 | Principal Substituent Chains in Isotopically Labeled Compounds |
| P-46.2.1 | Greater number of isotopically modified atoms |
| P-46.2.2 | Greater number of nuclides of higher mass number or isotopically modified substituent atoms or groups |
| P-46.3 | Principal Substituent Chains in Compounds with Stereogenic Centers |
| P-46.3.1 | Greater number of (<i>Z</i>)-double bonds |
| P-46.3.2 | Greater number of (<i>R</i>)-chirality centers |

P-5 SELECTING PREFERRED IUPAC NAMES AND CONSTRUCTING NAMES OF ORGANIC COMPOUNDS

| | |
|-------------|--|
| P-50 | Introduction |
| P-51 | Selecting the Preferred Type of IUPAC Nomenclature |
| P-51.0 | Introduction |
| P-51.1 | Selecting the Preferred Type of Nomenclature |
| P-51.1.1 | Substitutive nomenclature is preferred to functional class nomenclature |
| P-51.1.2 | Substitutive nomenclature is preferred to conjunctive nomenclature |
| P-51.1.3 | Skeletal replacement ('a') nomenclature is preferred to substitutive nomenclature |
| P-51.1.4 | Skeletal replacement ('a') nomenclature is preferred to multiplicative nomenclature |
| P-51.1.5 | Multiplicative nomenclature is preferred to simple substitutive nomenclature |
| P-51.2 | Functional Class Nomenclature |
| P-51.2.1 | Characteristic groups with preferred IUPAC names using functional class nomenclature |
| P-51.2.2 | Functional class nomenclature for general nomenclature |
| P-51.3 | Multiplicative Nomenclature |
| P-51.3.1 | Preferred IUPAC multiplicative names |
| P-51.3.2 | When more than two identical parent structures occur in the structure |

| | |
|-------------|---|
| P-51.3.3 | When conditions (1), (2), and (3) as defined in P-53.3.1 are not met |
| P-51.4 | Skeletal Replacement ('a') Nomenclature |
| P-51.4.1 | Skeletal replacement ('a') nomenclature in acyclic chains |
| P-51.4.2 | Skeletal replacement ('a') nomenclature for cyclic compounds |
| P-51.5 | Conjunctive Nomenclature <i>vs.</i> Substitutive Nomenclature |
| P-52 | Selecting Preferred IUPAC Names and Preselected Names for Parent Hydrides |
| P-52.1 | Selecting Preselected Names |
| P-52.1.1 | Mononuclear parent hydride names |
| P-52.1.2 | Preselected names for homogeneous acyclic polynuclear parent hydrides |
| P-52.1.3 | Preselected names for heterogeneous acyclic parent hydrides |
| P-52.1.4 | Preselected names for parent hydrides with nonstandard bonding numbers |
| P-52.1.5 | Heteromonocyclic noncarbon Hantzsch-Widman parent hydrides |
| P-52.1.6 | Heterocyclic noncarbon von Baeyer and spiro compounds |
| P-52.1.7 | Preselected names for homogeneous and heterogeneous heterobi- and heteropolycyclic fused ring systems |
| P-52.2 | Selecting Preferred IUPAC Names |
| P-52.2.1 | Acyclic and monocyclic hydrocarbons |
| P-52.2.2 | Heteroacyclic and heteromonocycles |
| P-52.2.3 | Unsaturated heteromonocyclic compounds with more than ten ring members |
| P-52.2.4 | Preferred IUPAC names in fusion nomenclature |
| P-52.2.4.1 | Five-membered ring requirement |
| P-52.2.4.2 | Heteromonocycles as components in fusion names |
| P-52.2.4.3 | Multiparent fused ring systems with three or more interparent components |
| P-52.2.4.4 | Limitations to fusion nomenclature |
| P-52.2.5 | Preferred IUPAC names in phane nomenclature |
| P-52.2.6 | Preferred IUPAC names for (C ₆₀ -I _n)[5,6]fullerene and (C ₇₀ -D _{5h(6)})[5,6]fullerene with 'nor' or 'seco' prefixes |
| P-52.2.7 | Preferred IUPAC names and numbering for ring assemblies |
| P-52.2.8 | Selection between a ring and a chain as parent hydride |
| P-53 | Selecting Preferred Retained Names of Parent Hydrides |
| P-54 | Selecting the Preferred Method for Modifying the Degree of Hydrogenation |
| P-54.1 | Methods for Modifying the Degree of Hydrogenation of Parent Hydrides: |
| P-54.2 | Unsaturated Monocyclic Carbocycles |
| P-54.3 | Unsaturation in Ring Assemblies Composed of Mancude and Saturated Rings |
| P-54.4 | Names Modified by 'hydro' or 'dehydro' Prefixes |
| P-54.4.1 | Hantzsch-Widman heteromonocycles |
| P-54.4.2 | Saturated heteromonocyclic compounds |
| P-54.4.3 | Fused ring systems and monocyclic mancude ring assemblies |

| | |
|-------------|---|
| P-54.4.4 | Names modified by 'dehydro' prefixes |
| P-55 | Selecting the Preferred Retained Name for Functional Parent Compounds |
| P-56 | Selecting the Preferred Suffix for the Principal Characteristic Group |
| P-56.1 | The Suffix 'peroxol' for -OOH |
| P-56.2 | The Suffixes 'SO-thioperoxol', and Chalogen Analogues |
| P-56.3 | The Suffixes 'imidamide' and 'carboximidamide' |
| P-56.4 | The Endings 'diyl' and 'ylidene' vs. 'ylene' |
| P-57 | Selecting Preferred and Preselected Prefixes for Substituent Groups |
| P-57.1 | Prefixes Derived from Parent Hydrides |
| P-57.1.1 | Prefixes derived from mononuclear and acyclic parent hydrides |
| P-57.1.2 | Retained names used as preferred prefixes |
| P-57.1.3 | Retained prefixes recommended only for general nomenclature |
| P-57.1.4 | Retained prefixes no longer recommended |
| P-57.1.5 | Prefixes derived from cyclic parent hydrides |
| P-57.1.6 | Prefixes derived from parent hydrides with modified degrees of hydrogenation |
| P-57.2 | Prefixes Derived from Characteristic (functional) Groups |
| P-57.3 | Prefixes Derived from Organic Functional Parent Compounds |
| P-57.4 | Construction of Linear Compound and/or Complex Substituent Prefixes. |
| P-58 | Selecting Preferred IUPAC Names |
| P-58.1 | Introduction |
| P-58.2 | Indicated Hydrogen, 'Added Indicated Hydrogen', and Nondetachable Hydro Prefixes |
| P-58.2.1 | Indicated hydrogen |
| P-58.2.2 | 'Added indicated hydrogen' |
| P-58.2.3 | Specific rules related to indicated hydrogen, 'added indicated hydrogen' and hydro prefixes |
| P-58.2.4 | Prefix nomenclature |
| P-58.2.5 | Nondetachable hydro prefixes vs. indicated hydrogen |
| P-58.3 | Homogeneous Hetero Chains and Functional Groups |
| P-58.3.1 | Preselected names for unbranched homogeneous heteroacyclic parent hydrides |
| P-58.3.2 | Atoms expressed by a principal characteristic group, a functional parent compound, or a compulsory prefix |
| P-59 | Name Construction |
| P-59.0 | Introduction |
| P-59.1 | General Methodology |
| P-59.1.1 | Determine the type(s) of nomenclature and operations to be used |
| P-59.1.2 | Determine the type(s) of nomenclature and operations to be used |
| P-59.1.3 | Select the parent hydride(s) |
| P-59.1.4 | Name the parent hydride, and the principal characteristic group |
| P-59.1.5 | Determine affixes and/or prefixes |

| | |
|-----------|--|
| P-59.1.6 | Name substituent groups and characteristic groups not cited as principal characteristic groups as prefixes |
| P-59.1.7 | Assemble the components into a complete name |
| P-59.1.8 | Complete the name with all required descriptors for stereochemical features |
| P-59.1.9 | Characteristic groups |
| P-59.1.10 | Numbering nomenclatural features |
| P-59.2 | Examples Illustrating the General Methodology |
| P-59.2.1 | Selection of parent hydrides |
| P-59.2.2 | Seniority of heteroatoms over suffixes |
| P-59.2.3 | Seniority of principal characteristic groups over unsaturation |
| P-59.2.4 | Seniority of 'ene' and 'yne' endings and hydro prefixes over detachable prefixes |
| P-59.2.5 | Treatment of detachable prefixes |

P-6 APPLICATIONS TO SPECIFIC CLASSES OF COMPOUNDS

| | |
|-------------|--|
| P-60 | Introduction |
| P-60.1 | Topical Outline |
| P-60.2 | Presentation of Names |
| P-61 | Substitutive Nomenclature: Prefix Mode |
| P-61.0 | Introduction |
| P-61.1 | General Methodology |
| P-61.2 | Hydrocarbyl Groups and Corresponding Di- and Polyvalent Groups |
| P-61.2.1 | Acyclic hydrocarbons |
| P-61.2.2 | Cyclic hydrocarbons |
| P-61.2.3 | Hydrocarbons consisting of rings and chains |
| P-61.2.4 | Structures containing heterocycles |
| P-61.3 | Halogen Compounds |
| P-61.3.1 | Halogen atoms in its standard bonding number |
| P-61.3.2 | Halogen atoms attached to heteroatoms |
| P-61.3.3 | Compounds containing the group $-I(OH)_2$ or similar groups |
| P-61.3.4 | Retained names |
| P-61.4 | Diazo Compounds |
| P-61.5 | Nitro and Nitroso Compounds |
| P-61.5.1 | Nitro and nitroso compounds |
| P-61.5.2 | Nitro attached to heteroatoms |
| P-61.5.3 | <i>aci</i> -Nitro compounds |
| P-61.6 | Heterones |
| P-61.7 | Azides |
| P-61.8 | Isocyanates |
| P-61.9 | Isocyanides |
| P-61.10 | Fulminates and Isofulminates |
| P-61.11 | Polyfunctional Compounds |

| | |
|-------------|---|
| P-61.11.1 | Low locants assigned as a set |
| P-61.11.2 | Low locants assigned to the prefix cited first in the name |
| P-62 | Amines and Imines |
| P-62.0 | Introduction |
| P-62.1 | General Methodology |
| P-62.2 | Amines |
| P-62.2.1 | Primary amines |
| P-62.2.2 | Secondary and tertiary amines |
| P-62.2.3 | The prefix 'amino' |
| P-62.2.4 | Polyamines |
| P-62.2.5 | Multiplicative nomenclature |
| P-62.2.6 | Modification of the degree of saturation/unsaturation of amines |
| P-62.3 | Imines |
| P-62.3.1 | Substitutive names for imines |
| P-62.4 | <i>N</i> -Substitution of Amines and Imines by Heteroatoms |
| P-62.5 | Amine Oxides, Imine Oxides, and Chalcogen Analogues |
| P-62.6 | Amine and Imine Salts |
| P-62.6.1 | Cation and anion names |
| P-62.6.2 | Salts of indefinite structure (adducts) |
| P-63 | Hydroxy Compounds, Ethers, Peroxols, Peroxides, and Chalcogen Analogues |
| P-63.0 | Introduction |
| P-63.1 | Hydroxy Compounds and Chalcogen Analogues |
| P-63.1.1 | Retained names |
| P-63.1.2 | Systematic names of alcohols, phenols, enols, and ynols |
| P-63.1.3 | Heterols |
| P-63.1.4 | Substitutive nomenclature, prefix mode |
| P-63.1.5 | Sulfur, selenium and tellurium analogues of hydroxy compounds |
| P-63.2 | Ethers and Chalcogen Analogues |
| P-63.2.1 | Definitions and general methodology |
| P-63.2.2 | Names of substituent groups R'-O-, R'-S-, R'-Se-, and R'-Te- |
| P-63.2.3 | Retained names of ethers |
| P-63.2.4 | Systematic names of ethers |
| P-63.2.5 | Names of chalcogen analogues of ethers, i.e., sulfides, selenides, and tellurides |
| P-63.3 | Peroxides and Chalcogen Analogues |
| P-63.3.1 | Peroxides, disulfides, diselenides, and ditellurides |
| P-63.3.2 | Mixed chalcogen analogues of peroxides |
| P-63.4 | Hydroperoxides (Peroxols) and Chalcogen Analogues |
| P-63.4.1 | Hydroperoxides |
| P-63.4.2 | Chalcogen analogues of hydroperoxides |
| P-63.5 | Cyclic Ethers, Sulfides, Selenides, and Tellurides |

| | |
|-------------|--|
| P-63.6 | Sulfoxides and Sulfones |
| P-63.7 | Polyfunctional Compounds |
| P-63.8 | Salts of Hydroxy Compounds, Hydroperoxy Compounds and Their Chalcogen Analogues |
| P-63.8.1 | Neutral salts of hydroxy compounds and their chalcogen analogues |
| P-63.8.2 | Cyclic salts are named as heterocycles |
| P-63.8.3 | Partial salts of polyols and their chalcogen |
| P-64 | Ketones, Pseudoketones, Heterones, and Chalcogen Analogues |
| P-64.0 | Introduction |
| P-64.1 | Definitions |
| P-64.1.1 | Ketones |
| P-64.1.2 | Pseudoketones and heterones |
| P-64.2 | Ketones |
| P-64.2.1 | Retained names |
| P-64.2.2 | Systematic construction of names for ketones |
| P-64.2.2.1 | Acyclic ketones |
| P-64.2.2.2 | Cyclic ketones |
| P-64.2.2.3 | Seniority order for numbering |
| P-64.2.2.4 | Ketenes |
| P-64.3 | Pseudoketones |
| P-64.3.1 | Cyclic anhydrides, esters and amides |
| P-64.3.2 | Acyclic pseudoketones |
| P-64.4 | Heterones |
| P-64.4.1 | Acyclic heterones |
| P-64.4.2 | Thioketone and thioaldehyde oxides |
| P-64.5 | Carbonyl Groups as Prefixes |
| P-64.5.1 | Ketones |
| P-64.5.2 | Pseudoketones |
| P-64.6 | Chalcogen Analogues of Ketones, Pseudoketones, and Heterones |
| P-64.6.1 | Chalcogen analogues of ketones, pseudoketones and heterones |
| P-64.6.2 | Seniority order of suffixes |
| P-64.7 | Polyfunctional Ketones, Pseudoketones and Heterones |
| P-64.7.1 | Ketones, pseudoketones and heterones, and their chalcogen analogues |
| P-64.7.2 | There is no seniority order difference between ketones and pseudoketones |
| P-64.7.3 | When an oxygen atom of a ketone, pseudoketone, or heterone is replaced by a sulfur, selenium or tellurium atom |
| P-64.8 | Acyloins |
| P-65 | Acids, Acyl Halides and Pseudohalides, Salts, Esters, and Anhydrides |
| P-65.0 | Introduction |
| P-65.1 | Carboxylic Acids and Functional Replacement Analogues |
| P-65.1.1 | Retained names |
| P-65.1.2 | Systematic names |

| | |
|------------|--|
| P-65.1.2.1 | Carboxylic acid groups |
| P-65.1.2.2 | The suffix 'carboxylic acid' |
| P-65.1.2.3 | Seniority order for numbering |
| P-65.1.2.4 | Polyfunctional carboxylic acids |
| P-65.1.3 | Carboximidic, carbohydrazonic, carbohydroximic, and carbohydroxamic acids |
| P-65.1.3.1 | Carboximidic acids |
| P-65.1.3.2 | Carbohydrazonic acids |
| P-65.1.3.3 | Carbohydroximic acids |
| P-65.1.4 | Peroxy-carboxylic acids |
| P-65.1.5 | Chalcogen analogues of carboxylic acids |
| P-65.1.5.1 | Functional replacement in systematic names of carboxylic acids |
| P-65.1.5.2 | Functional replacement in retained names of carboxylic acids |
| P-65.1.5.3 | Functional replacement in peroxy-carboxylic acids |
| P-65.1.6 | Amic, anilic, and aldehydic acids |
| P-65.1.7 | Acyl groups derived from carboxylic and related acids |
| P-65.1.7.1 | Definitions and name formation |
| P-65.1.7.2 | Carboxylic acids having retained names that are preferred IUPAC names |
| P-65.1.7.3 | Carboxylic acids with names retained only for general nomenclature |
| P-65.1.7.4 | Systematically named carboxylic acids |
| P-65.1.7.5 | Mixed acyl groups |
| P-65.1.8 | Formic acid |
| P-65.2 | Carbonic, Cyanic, and Di- and Polycarbonic Acids |
| P-65.2.1 | Carbonic acid |
| P-65.2.2 | Cyanic acid |
| P-65.2.3 | Di-, tri-, tetra-, and polycarbonic acids |
| P-65.3 | Sulfur, Selenium, Tellurium Acids with Chalcogen Atoms Directly Linked to a Parent Hydride |
| P-65.3.0 | Introduction |
| P-65.3.1 | Substitutive nomenclature, suffix mode, for sulfonic, sulfinic, etc., acid |
| P-65.3.2 | Substitutive nomenclature, prefix mode for sulfonic, sulfinic, etc. acids |
| P-65.3.3 | Polyfunctional compounds |
| P-65.4 | Acyl Groups as Substituent Groups |
| P-65.4.1 | General methodology |
| P-65.5 | Acyl Halides and Pseudohalides |
| P-65.5.1 | Acyl halides from suffix acids |
| P-65.5.2 | Acyl pseudohalides from suffix acids |
| P-65.5.3 | Acyl halides and pseudohalides from carbonic, cyanic, and the polycarbonic acids |
| P-65.5.4 | Acyl halides and pseudohalides as substituent groups |
| P-65.6 | Salts and Esters |
| P-65.6.1 | General methodology |

| | |
|--------------|---|
| P-65.6.2 | Salts |
| P-65.6.3 | Esters, lactones, and related compounds |
| P-65.6.3.1 | Definitions |
| P-65.6.3.2 | General methodology |
| P-65.6.3.3 | Preferred IUPAC names for esters |
| P-65.6.3.3.1 | Monoesters |
| P-65.6.3.3.2 | Polyesters derived from a single acid component |
| P-65.6.3.3.3 | Polyesters formed from a single 'alcoholic' component |
| P-65.6.3.3.4 | Polyesters derived from multiple acid components and multiple 'alcoholic' components |
| P-65.6.3.3.5 | Partial esters from polybasic acids and salts |
| P-65.6.3.3.6 | Substitutive nomenclature is senior to functional class nomenclature for preferred IUPAC names for esters |
| P-65.6.3.3.7 | Esters of acids modified by functional replacement |
| P-65.6.3.4 | Pseudoesters |
| P-65.6.3.5 | Cyclic esters |
| P-65.6.3.6 | Acylals |
| P-65.7 | Anhydrides and Their Analogues |
| P-65.7.0 | Introduction |
| P-65.7.1 | Symmetric anhydrides |
| P-65.7.2 | Mixed anhydrides |
| P-65.7.3 | Thioanhydrides and other chalcogen analogues |
| P-65.7.4 | Peroxyanhydrides and chalcogen analogues |
| P-65.7.5 | Diacyl derivatives of trioxidane and chalcogen analogues |
| P-65.7.6 | Di- and polyanhydrides |
| P-65.7.7 | Cyclic anhydrides |
| P-65.7.8 | Polyfunctional anhydrides |
| P-66 | Amides, Imides, Hydrazides, Nitriles, and Aldehydes |
| P-66.0 | Introduction |
| P-66.1 | Amides |
| P-66.1.0 | Introduction |
| P-66.1.1 | Primary amides |
| P-66.1.1.1 | Carboxamides |
| P-66.1.1.2 | Sulfonamides, sulfinamides, and related selenium and tellurium amides |
| P-66.1.1.3 | Substitution of primary amides |
| P-66.1.1.4 | Amides denoted as prefixes |
| P-66.1.2 | Secondary and tertiary amides |
| P-66.1.3 | 'Hidden' amides |
| P-66.1.4 | Chalcogen analogues of amides |
| P-66.1.5 | Lactams, lactims, sultams, and sultims |
| P-66.1.6 | Amides derived from carbonic, cyanic, and the di- and polycarbonic acids |

| | |
|------------|---|
| P-66.1.6.1 | Carbonic acid and related compounds |
| P-66.1.6.2 | Cyanic acid |
| P-66.1.6.3 | Di- and polycarbonic acids |
| P-66.1.7 | Polyfunctional amides |
| P-66.2 | Imides |
| P-66.2.1 | Acyclic imides and cyclic imides |
| P-66.2.2 | Imide prefixes |
| P-66.3 | Hydrazides |
| P-66.3.0 | Definition |
| P-66.3.1 | Systematic names |
| P-66.3.2 | Substituent groups derived from hydrazides |
| P-66.3.3 | Substituted hydrazides |
| P-66.3.4 | Chalogen analogues of hydrazides |
| P-66.3.5 | Hydrazides from carbonic, cyanic, and di- and the polycarbonic acids |
| P-66.3.6 | Semioxamazones |
| P-66.4 | Amidines, Amidrazones, Hydrazidines, and Amidoximes (Amide Oximes) |
| P-66.4.1 | Amidines |
| P-66.4.1.1 | Suffixes for amidines |
| P-66.4.1.2 | Amidines of carbonic, and di- and polycarbonic acids |
| P-66.4.1.3 | Prefixes for the amidine characteristic group |
| P-66.4.1.4 | Substituted amidines |
| P-66.4.1.5 | Formamidine disulfides |
| P-66.4.1.6 | Diamidides |
| P-66.4.2 | Amidrazones |
| P-66.4.2.1 | Amidrazone suffixes |
| P-66.4.2.2 | Amidrazones of carbonic acid and di- and polycarbonic acids |
| P-66.4.2.3 | Amidrazone prefixes |
| P-66.4.3 | Hydrazidines |
| P-66.4.3.1 | Hydrazidine suffixes |
| P-66.4.3.2 | Hydrazidines derived from sulfonic acid, and similar selenium and tellurium acids |
| P-66.4.3.3 | Hydrazidines derived from carbonic acid and di- and polycarbonic acids |
| P-66.4.3.4 | Hydrazidine prefixes |
| P-66.4.4 | Amidoximes (amide oximes) |
| P-66.5 | Nitriles |
| P-66.5.0 | Introduction |
| P-66.5.1 | Preferred names for nitriles |
| P-66.5.2 | Substituted nitriles |
| P-66.5.3 | Nitriles/cyanides corresponding to carbonic and di- and polycarbonic acids |
| P-66.5.4 | Nitrile oxides and chalcogen analogues |
| P-66.6 | Aldehydes |

| | |
|--------------|--|
| P-66.6.0 | Introduction |
| P-66.6.1 | Systematic names of aldehydes |
| P-66.6.2 | Aldehydes from di- and polycarbonic acids |
| P-66.6.3 | Chalcogen analogues of aldehydes |
| P-66.6.4 | Polyfunctional aldehydes |
| P-66.6.5 | Acetals and ketals, hemiacetals and hemiketals, and their chalcogen analogues |
| P-67 | Mononuclear and Polynuclear Noncarbon Acids and their Functional Replacement Analogues as Functional Parents for Naming Organic Compounds |
| P-67.0 | Introduction |
| P-67.1 | Mononuclear Noncarbon Oxoacids |
| P-67.1.1 | Names of mononuclear noncarbon oxoacids |
| P-67.1.2 | Functional replacement nomenclature applied to mononuclear noncarbon oxoacids |
| P-67.1.2.1 | Mononuclear acids modified by infixes |
| P-67.1.2.2 | Mononuclear acids modified by prefixes |
| P-67.1.2.3 | General methodology for functional replacement nomenclature using infixes |
| P-67.1.2.4 | Mononuclear noncarbon oxoacids modified by functional replacement nomenclature |
| P-67.1.2.5 | Acid halides and pseudohalides |
| P-67.1.2.6 | Amides and hydrazides |
| P-67.1.3 | Salts, esters and anhydrides of mononuclear noncarbon oxoacids |
| P-67.1.4 | Substituent prefix groups derived from mononuclear noncarbon oxoacids |
| P-67.1.4.1 | Mononuclear nitrogen, phosphorus, arsenic, and antimony acids |
| P-67.1.4.1.1 | Preselected prefixes |
| P-67.1.4.1.2 | Substituent groups for general nomenclature |
| P-67.1.4.1.3 | Compound and complex substituent groups |
| P-67.1.4.2 | Substituent groups derived from boron acid and silicic acid |
| P-67.1.4.3 | Substituent groups derived from nitric and nitrous acids |
| P-67.1.4.4 | Substituent groups derived from chalcogen acids |
| P-67.1.4.5 | Substituent groups derived from halogen acids |
| P-67.1.5 | Seniority order among noncarbon oxoacids and derivatives |
| P-67.1.6 | <i>aci</i> -Nitro compounds |
| P-67.2 | Di- and Polynuclear Noncarbon Oxoacids |
| P-67.2.1 | Preselected names |
| P-67.2.2 | Functional replacement derivatives of di- and polynuclear noncarbon oxoacids |
| P-67.2.3 | Acid halides and pseudohalides of di- and polynuclear noncarbon oxoacids |
| P-67.2.4 | Amides and hydrazides of di- and polynuclear noncarbon oxoacids |
| P-67.2.5 | Esters and anhydrides of di- and polynuclear noncarbon oxoacids |

| | |
|--------------|--|
| P-67.2.6 | Substituent groups derived from polyacids |
| P-67.3 | Substitutive Names and Functional Class Names of Polyacids |
| P-67.3.1 | Polynuclear noncarbon oxoacids that cannot be formed on the basis of fundamental oxoacids |
| P-67.3.2 | The disulfurous acid name dilemma |
| P-68 | Nomenclature of Organic Compounds of the Group 13, 14, 15, 16, and 17 Elements not Included in Sections P-62 through P-67 |
| P-68.0 | Introduction |
| P-68.1 | Nomenclature of Compounds of Group 13 Elements |
| P-68.1.1 | Parent hydrides |
| P-68.1.2 | Names for substituent groups derived from parent hydrides |
| P-68.1.3 | Modification of the degree of hydrogenation |
| P-68.1.4 | Functional parent compounds |
| P-68.1.5 | Substitutive nomenclature |
| P-68.1.5.1 | Suffix nomenclature |
| P-68.1.5.2 | Prefix nomenclature |
| P-68.1.6 | Adducts |
| P-68.1.6.1 | Structures for Lewis 'adducts' |
| P-68.1.6.2 | General organic method for naming Lewis adducts |
| P-68.1.6.3 | Intramolecular adducts |
| P-68.2 | Nomenclature for Compounds of Group 14 Elements |
| P-68.2.0 | Introduction |
| P-68.2.1 | Silicon, germanium, tin, and lead parent hydrides |
| P-68.2.2 | Substituent groups derived from parent hydrides |
| P-68.2.3 | Modification of the degree of hydrogenation |
| P-68.2.4 | Silicic acid as a functional parent compound |
| P-68.2.5 | Substitutive nomenclature: suffix mode |
| P-68.2.6 | Substitutive nomenclature: prefix mode |
| P-68.3 | Nomenclature of Compounds of Group 15 Elements |
| P-68.3.1 | Nitrogen compounds |
| P-68.3.1.0 | Introduction |
| P-68.3.1.1 | Hydroxylamines, oximes, and nitrolic acids and nitrosolic acids |
| P-68.3.1.2 | Hydrazine and related compounds: hydrazones, azines, semicarbazides, semicarbazones, and carbonohydrazides |
| P-68.3.1.3 | Diazene and related compounds |
| P-68.3.1.3.1 | Substitution of diazene |
| P-68.3.1.3.2 | Azo compounds, R-N=N-R' |
| P-68.3.1.3.3 | Azoxy compounds, R-N=N(O)-R' |
| P-68.3.1.3.4 | Diazene-carbohydrazide, HN=N-CO-NH-NH ₂ |
| P-68.3.1.3.5 | Formazan, H ₂ N-N=CH-N=NH |
| P-68.3.1.3.6 | Carbodiazone [bis(diazenyl)methanone], HN=N-CO-N=NH |
| P-68.3.1.3.7 | Isodiazene, R ₂ N ⁺ =N ⁻ |
| P-68.3.1.4 | Polyazanes |

| | |
|-------------|--|
| P-68.3.2 | Phosphorus, arsenic, and antimony compounds |
| P-68.3.2.1 | General methodology |
| P-68.3.2.2 | Parent hydrides |
| P-68.3.2.3 | Substitutive nomenclature |
| P-68.3.3 | Bismuth compounds |
| P-68.4 | Nomenclature of Compounds of Group 16 Elements |
| P-68.4.0 | Introduction |
| P-68.4.1 | Three or more homogeneous contiguous chalcogen atoms |
| P-68.4.2 | Three or more heterogeneous contiguous chalcogen atoms |
| P-68.4.3 | Chalcogen parent hydrides with nonstandard bonding numbers |
| P-68.4.3.1 | Sulfanes, selanes, and tellanes |
| P-68.4.3.2 | Di- and polysulfoxides, polysulfones, and selenium and tellurium analogues |
| P-68.4.3.3 | Sulfimides, and chalcogen analogues, $H_2E=NH$, where E = S, Se, or Te |
| P-68.4.3.4 | Sulfinylamines, $RN=E=O$, sulfonylamines, $RN=E(=O)_2$ and chalcogen analogues where E = S, Se, or Te |
| P-68.4.3.5 | Sulfonediimines, $RE(=NH)_2R'$ and chalcogen compounds where E = S, Se, or Te |
| P-68.4.3.6 | Sulfoximides, $R_2E(=O)=NR'$ and chalcogen analogues where E = S, Se, or Te |
| P-68.4.3.7 | Sulfur diimides, $HN=E=NH$ and chalcogen analogues where E = S, Se, or Te |
| P-68.4.3.8 | Sulfur triimides, $E(=NH)_3$ and chalcogen analogues where E = S, Se, or Te |
| P-68.5 | Nomenclature of Compounds of Group 17 Elements |
| P-68.5.0 | Substitutive nomenclature |
| P-68.5.1 | Nomenclature based on halogen parent hydrides |
| P-68.5.2 | Nomenclature of halogen acids |
| P-68.5.3 | Amides of halogen acids |
| P-69 | Nomenclature for Organometallic Compounds |
| P-69.0 | Introduction |
| P-69.1 | Organometallic Compounds of Elements in Groups 13, 14, 15, and 16 |
| P-69.2 | Organometallic Compounds of Elements in Groups 3 through 12 |
| P-69.2.1 | Coordination nomenclature |
| P-69.2.2 | Definitions of terms |
| P-69.2.3 | Compounds with at least one metal-carbon single bond |
| P-69.2.4 | Organometallic groups with multicenter bonding to carbon atoms |
| P-69.2.5 | Bridging organometallic groups with multicenter bonding to carbon atoms |
| P-69.2.6 | Organometallic compounds with unsaturated molecules and substituent groups |
| P-69.2.7 | “Ocenes” |
| P-69.3 | Organometallic Compounds of Elements in Groups 1 and 2 |

| | |
|----------|--|
| P-69.4 | Metallacycles |
| P-69.5 | Order of Seniority Order for Organometallic Compounds |
| P-69.5.1 | Compounds having two identical or different metal atoms |
| P-69.5.2 | Compounds having one metal atom of class (1) and another atom of class (2) |
| P-69.5.3 | For organometallic compounds having two metal atoms belonging to class (2) |

P-7 RADICALS, IONS, AND RELATED SPECIES

| | |
|-------------|--|
| P-70 | Introduction |
| P-70.1 | General Methodology |
| P-70.2 | Seniority of Radicals and Ions |
| P-70.3 | Name Formation |
| P-70.3.1 | Suffixes, prefixes, and endings for radicals and ions in substitutive nomenclature |
| P-70.3.2 | Basic multiplying prefixes |
| P-70.3.3 | Order of citing suffixes and endings |
| P-70.4 | General Rules for the Selection of Preferred Names |
| P-71 | Radicals |
| P-71.1 | General Methodology |
| P-71.2 | Radicals Derived from Parent Hydrides |
| P-71.2.1 | Monovalent radicals |
| P-71.2.2 | Divalent and trivalent radicals |
| P-71.2.3 | Multiple radical centers (polyradicals) |
| P-71.2.4 | Acyclic radicals derived by the removal of one or more hydrogen atoms from nonterminal chain positions |
| P-71.2.5 | The λ -convention |
| P-71.2.6 | ‘Added indicated hydrogen’ for radicals of mancude systems |
| P-71.3 | Radical Centers on Characteristic Groups |
| P-71.3.1 | Acyl radicals |
| P-71.3.2 | Removal of hydrogen atoms from an amine, imine, or amide characteristic group |
| P-71.3.3 | Polyamine, polyimine and polyamide radicals |
| P-71.3.4 | Removal of the hydrogen atom of a hydroxy group (or chalcogen analogue) of an acid or hydroxy compound |
| P-71.4 | Assemblies of Parent Radicals |
| P-71.5 | Prefixes Denoting Radicals |
| P-71.6 | Order of Citation and Seniority of Suffixes ‘yl’, ‘ylidene’, and ‘ylidyne’ |
| P-71.7 | Choice of Parent Radical |
| P-72 | Anions |
| P-72.1 | General Methodology |
| P-72.2 | Anions Formed by Removal of Hydrons |
| P-72.2.1 | Functional class nomenclature |
| P-72.2.2 | Systematic nomenclature |

| | |
|--------------|---|
| P-72.2.2.1 | Anions derived from parent hydrides and their derivatives |
| P-72.2.2.2 | Anions derived from characteristic groups |
| P-72.2.2.2.1 | Anions derived from acids |
| P-72.2.2.2.2 | Anions derived from hydroxy compounds |
| P-72.2.2.2.3 | Anions derived from amines and imines |
| P-72.2.2.2.4 | Anions derived from other characteristic groups |
| P-72.3 | Anions Formed by Addition of Hydride Ions |
| P-72.4 | Skeletal Replacement ('a') Nomenclature |
| P-72.5 | Multiple Anionic Centers |
| P-72.5.1 | Assemblies of parent anions |
| P-72.5.2 | 'Ide' and 'uide' centers in the same parent hydride |
| P-72.5.3 | Anionic characteristic groups on anionic parent hydrides |
| P-72.6 | Anionic Centers in Both Parent Compounds and Substituent Groups |
| P-72.6.1 | Prefixes for anionic centers derived from acid characteristic groups |
| P-72.6.2 | Prefixes for anionic chalcogens |
| P-72.6.3 | Systematically formed prefixes that include anionic center(s) |
| P-72.7 | Choice of an Anionic Parent Structure |
| P-72.8 | The Suffixes 'ide' and 'uide' and the λ -Convention |
| P-72.8.1 | The suffix 'uide' is preferred to the suffix 'ide' with a parent hydride named by the λ -convention |
| P-72.8.2 | The suffix 'ide' is preferred to the suffix 'uide' involved with the λ -convention |
| P-73 | Cations |
| P-73.0 | Introduction |
| P-73.1 | Cationic Compounds with Cationic Centers Derived Formally by the Addition of Hydrons |
| P-73.1.1 | Cationic centers in parent hydrides |
| P-73.1.2 | Cationic centers on characteristic groups |
| P-73.1.2.1 | Cationic compounds derived from neutral compounds expressed by suffixes |
| P-73.1.2.2 | Uronium ions and chalcogen analogues |
| P-73.2 | Cationic Compounds with Cationic Centers Derived Formally by the Removal of Hydride Ions |
| P-73.2.1 | Functional class names |
| P-73.2.2 | Cationic centers in parent hydrides |
| P-73.2.2.1 | Cationic centers in parent hydrides |
| P-73.2.2.2 | 'Added indicated hydrogen' |
| P-73.2.2.3 | Diazonium ions |
| P-73.2.3 | Cationic centers on characteristic groups |
| P-73.3 | The λ -Convention with the Suffix 'ylium' |
| P-73.3.1 | Application of the λ -convention with the suffix 'ylium' |
| P-73.3.2 | Retained names |
| P-73.4 | Skeletal Replacement ('a') Nomenclature for Cations |

| | |
|-------------|--|
| P-73.5 | Cationic Compounds with Multiple Cationic Centers |
| P-73.5.1 | Assemblies of parent cations |
| P-73.5.2 | 'Ium' and 'ylium' centers in the same parent hydride |
| P-73.5.3 | Cationic characteristic groups on parent cations |
| P-73.6 | Cationic Prefix Names |
| P-73.7 | Choice of a Parent Structure |
| P-73.8 | The Suffixes 'ium' vs. 'ylium' and the λ -convention |
| P-73.8.1 | The suffix '-ylium' is preferred to the suffix '-ium' with a parent hydride modified by the λ -convention |
| P-73.8.2 | The suffix 'ium' is preferred to the suffix 'ylium' with a parent hydride modified by the λ -convention |
| P-74 | Zwitterions |
| P-74.0 | Introduction |
| P-74.1 | Zwitterionic Parent Structures having the Anionic and Cationic centers on the same Parent Compound Including Ionic Centers on Characteristic Groups Expressible as Suffixes. |
| P-74.1.1 | Ionic centers in the same parent structure |
| P-74.1.2 | Zwitterionic compounds with at least one ionic center on a characteristic group |
| P-74.1.3 | Anionic and cationic centers on different parent structures |
| P-74.2 | Dipolar Ions |
| P-74.2.1 | 1,2-Dipolar compounds |
| P-74.2.1.1 | 'Ylides' |
| P-74.2.1.2 | Amine oxides, imine oxides, and their chalcogen analogues |
| P-74.2.1.3 | Amine imides |
| P-74.2.1.4 | Phosphine oxides and chalcogen analogues |
| P-74.2.1.5 | Phosphine imides |
| P-74.2.2 | 1,3-Dipolar compounds |
| P-74.2.2.1 | Allyl type compounds with delocalized general structure |
| P-74.2.2.2 | The propargyl (propynyl) type including compounds with canonical resonance forms |
| P-74.2.2.3 | The carbene type including compounds having with canonical resonance forms |
| P-74.2.3 | Dipolar substituent groups |
| P-75 | Radical Ions |
| P-75.1 | Radical Ions Formed by the Addition or Removal of of Electrons |
| P-75.2 | Radical Ions Derived from Parent Hydrides |
| P-75.2.1 | Examples of radical anions |
| P-75.2.2 | Examples of radical cations |
| P-75.2.3 | Examples of zwitterionic radical ions |
| P-75.2.4 | 'Added indicated hydrogen' |
| P-75.3 | Radical Ions on Characteristic Groups |
| P-75.3.1 | Radical ions on ionic suffix groups |
| P-75.3.2 | Radical ions other than those named by using ionic suffix groups |

| | |
|-------------|--|
| P-75.4 | Ionic and Radical Centers in Different Parent Structures |
| P-76 | Delocalized Radicals and Ions |
| P-76.1 | Delocalization in Names Involving One Radical or Ionic Center in an Otherwise Conjugated Double Bonds Structure is Denoted by the Appropriate Suffix without Locants |
| P-77 | Salts |
| P-77.1 | Preferred Names for Salts of Organic Bases |
| P-77.1.1 | Preferred IUPAC names for salts of organic bases are binary names |
| P-77.1.2 | Substitutive nomenclature is used to derive preferred IUPAC names for monosalts of di- or polyamines |
| P-77.1.3 | Traditional methods of naming salts of organic base when P-77.1.2 cannot be applied |
| P-77.2 | Salts Derived from Alcohols (including Phenols), Peroxols, and their Chalcogen Analogues |
| P-77.2.1 | Preferred IUPAC names are binary names formed by citing the name of the cation followed by that of the anion |
| P-77.2.2 | Substitutive nomenclature is used to derive IUPAC preferred names for monosalts of polyhydroxy compounds |
| P-77.3 | Salts Derived from Organic Acids |
| P-77.3.1 | Preferred IUPAC names are binary names formed by citing the name of the cation followed by that of the anion |
| P-77.3.2 | Substitutive nomenclature is used to derive preferred IUPAC names for acid salts of polybasic organic acids |

P-8 ISOTOPICALLY MODIFIED COMPOUNDS

| | |
|-------------|---|
| P-80 | Introduction |
| P-81 | Symbols and Definitions |
| P-81.1 | Nuclide Symbols |
| P-81.2 | Atomic Symbols |
| P-81.3 | Names for Hydrogen Atoms and Ions |
| P-81.4 | Isotopically Unmodified Compounds |
| P-81.5 | Isotopically Modified Compounds |
| P-82 | Isotopically Substituted Compounds |
| P-82.0 | Introduction |
| P-82.1 | Formulas |
| P-82.2 | Names |
| P-82.2.1 | Isotopically substituted compounds |
| P-82.2.2 | Isotopically modified compounds |
| P-82.2.3 | Addition of hydro prefixes |
| P-82.2.4 | Location of isotopic descriptor in a two or more word name |
| P-82.2.5 | Location of isotopic descriptor in a one word name |
| P-82.3 | Order of Nuclide Symbols |
| P-82.3.1 | When isotopes of different elements are present as nuclides |
| P-82.3.2 | When several isotopes of the same element are present as nuclides |

| | |
|-------------|--|
| P-82.4 | Stereoisomeric Isotopically Substituted Compounds |
| P-82.4.1 | When stereoisomerism results from isotopic substitution |
| P-82.4.2 | Isotopically substituted stereoisomers |
| P-82.5 | Numbering |
| P-82.5.1 | Numbering in relation to the unmodified compound |
| P-82.5.2 | Priority between isotopically substituted and unmodified atoms or groups |
| P-82.6 | Locants |
| P-82.6.1 | Omission of locants |
| P-82.6.2 | Letter and/or numeral locants |
| P-82.6.3 | Location of nuclides on positions not normally denoted by locants |
| P-82.6.4 | Italicized nuclide symbols and/or italic capital letters to distinguish between different nuclides of the same element |
| P-83 | Isotopically Labeled Compounds |
| P-83.1 | Specifically Labeled Compounds |
| P-83.1.1 | Structures of specifically labeled compounds |
| P-83.1.2 | Names of specifically labeled compounds |
| P-83.2 | Selectively Labeled Compounds |
| P-83.2.1 | Structures of selectively labeled compounds |
| P-83.2.2 | Names of selectively labeled compounds |
| P-83.3 | Nonselectively Labeled Compounds |
| P-83.3.1 | Nonselectively labeled compounds |
| P-83.3.2 | Structures |
| P-83.3.3 | Names |
| P-83.4 | Isotopically Deficient Compounds |
| P-83.4.1 | Isotopically deficient compounds |
| P-83.4.2 | Structures |
| P-83.4.3 | Names |
| P-83.5 | General and Uniform Labeling |
| P-83.5.1 | Selectively labeled compound, all positions labeled, not necessarily evenly; the symbol 'G' |
| P-83.5.2 | Selectively labeled compound, all positions uniformly labeled; the symbol 'U' |
| P-83.5.3 | Selectively labeled compound, uniformly labelled at specified positions |
| P-84 | Comparative Examples of Formulas and Names of Isotopically Modified Compounds |

P-9 SPECIFICATION OF CONFIGURATION AND CONFORMATION

| | |
|-------------|---|
| P-90 | Introduction |
| P-91 | Stereoisomer Graphical Representation and Naming |
| P-91.1 | Stereoisomer Graphical Representation |
| P-91.2 | Stereodescriptors |

| | |
|-------------|---|
| P-91.2.1 | Recommended stereodescriptors |
| P-91.2.2 | Omission of stereodescriptors |
| P-91.3 | Naming of Stereoisomer |
| P-92 | CIP Priority and Sequence Rules |
| P-92.1 | The Cahn-Ingold-Prelog (CIP) System: General Methodology |
| P-92.1.1 | Stereogenic units |
| P-92.1.2 | Rules for the assignment of configuration |
| P-92.1.3 | The Sequence Rules |
| P-92.1.4 | Hierarchical digraphs |
| P-92.1.4.1 | Acyclic molecules |
| P-92.1.4.2 | Double and triple bonds |
| P-92.1.4.3 | Saturated rings and ring systems |
| P-92.1.4.4 | Mancude rings and ring systems |
| P-92.1.5 | Exploration of a hierarchical digraph |
| P-92.1.6 | Ligand ranking: application of the five Sequence Rules |
| P-92.2 | Sequence Rule 1 |
| P-92.2.1 | Sequence subrule 1a : Ligands are arranged in order of decreasing atomic number |
| P-92.2.1.1 | Saturated compounds |
| P-92.2.1.2 | Double and triple bonds |
| P-92.2.1.3 | Saturated rings and ring systems |
| P-92.2.2 | Sequence Subrule 1b : Priority due to duplicate atoms |
| P-92.3 | Sequence Rule 2: Higher atomic mass number precedes lower atomic mass number |
| P-92.4 | Sequence Rule 3 |
| P-92.4.1 | ‘ <i>seqcis</i> ’ = (‘ <i>Z</i> ’) and ‘ <i>seqtrans</i> ’ = (‘ <i>E</i> ’) |
| P-92.4.2 | Sequence Rule 3: ‘ <i>seqcis</i> ’ (‘ <i>Z</i> ’) precedes ‘ <i>seqtrans</i> ’ (‘ <i>E</i> ’) and this order precedes nonstereogenic double bonds |
| P-92.5 | Sequence Rule 4 |
| P-92.5.1 | Sequence Rule 4a : Chiral stereogenic units precede pseudoasymmetric and these precede nonstereogenic units |
| P-92.5.2 | Sequence Rule 4b : Like descriptor pairs have priority over unlike descriptor pairs |
| P-92.5.3 | Sequence rule 4c : ‘ <i>r</i> ’ precedes ‘ <i>s</i> ’ |
| P-92.6 | Sequence Rule 5: ‘ <i>R</i> ’ precedes ‘ <i>S</i> ’, ‘ <i>M</i> ’ precedes ‘ <i>P</i> ’, and ‘ <i>seqCis</i> ’ precedes ‘ <i>seqTrans</i> ’ |
| P-93 | Configuration Specification |
| P-93.0 | Introduction |
| P-93.1 | General Aspects of Configuration Specification |
| P-93.1.1 | Absolute configuration |
| P-93.1.2 | Relative configuration |
| P-93.1.3 | Racemates |
| P-93.1.4 | The descriptor ‘ <i>ambo</i> ’ |

| | |
|------------|---|
| P-93.2 | Tetrahedral Configuration of Atoms other than Carbon |
| P-93.2.0 | Introduction |
| P-93.2.1 | General methodology |
| P-93.2.2 | Azanium (ammonium) and phosphanium (phosphonium) salts |
| P-93.2.3 | Amine oxides and phosphane oxides |
| P-93.2.4 | Phosphates, phosphonates, and related compounds |
| P-93.2.5 | Sulfates, sulfonates, and related compounds |
| P-93.2.6 | Compounds of Group 14 (other than carbon compounds) |
| P-93.3 | Nontetrahedral Configuration |
| P-93.3.1 | General methodology |
| P-93.3.2 | Polyhedral symbols |
| P-93.3.3 | Configuration index and priority number |
| P-93.3.3.1 | General methodology |
| P-93.3.3.2 | Trigonal pyramid |
| P-93.3.3.3 | T-shape systems |
| P-93.3.3.4 | See-saw systems |
| P-93.3.3.5 | Trigonal bipyramid |
| P-93.3.3.6 | Square pyramid |
| P-93.3.3.7 | Octahedron |
| P-93.3.4 | Chirality symbols |
| P-93.3.4.1 | The chirality symbols ' <i>R/S</i> ' |
| P-93.3.4.2 | The chirality symbols ' <i>A</i> ' and ' <i>C</i> ' |
| P-93.4 | Configuration Specification of Acyclic Organic Compounds |
| P-93.4.1 | Specification of stereogenic centers |
| P-93.4.2 | Specification of configuration for double bonds |
| P-93.5 | Configuration Specification of Cyclic Organic Compounds |
| P-93.5.0 | Introduction |
| P-93.5.1 | Monocyclic compounds |
| P-93.5.1.1 | Specification of stereogenic centers: stereodescriptors ' <i>R</i> ', ' <i>S</i> ', ' <i>r</i> ', and ' <i>s</i> ' |
| P-93.5.1.2 | Relative configuration: stereodescriptors ' <i>cis</i> ' and ' <i>trans</i> ' |
| P-93.5.1.3 | Relative configuration: stereodescriptors ' <i>r</i> ', ' <i>c</i> ', and ' <i>t</i> ' |
| P-93.5.1.4 | Unsaturated alicyclic compounds |
| P-93.5.2 | von Baeyer compounds |
| P-93.5.2.1 | Specification of stereogenic centers by CIP stereodescriptors |
| P-93.5.2.2 | Relative configuration (' <i>endo</i> ', ' <i>exo</i> ', ' <i>syn</i> ', ' <i>anti</i> ', ' <i>cis</i> ', ' <i>trans</i> ') |
| P-93.5.2.3 | Specification of double bonds |
| P-93.5.3 | Spiro compounds |
| P-93.5.3.1 | Specification of stereogenic spiro atoms of the type ' <i>Xabcd</i> ', here ' <i>a</i> ' > ' <i>b</i> ' > ' <i>c</i> ' > ' <i>d</i> ' |
| P-93.5.3.2 | Specification of stereogenic spiro atoms of the type ' <i>Xabab</i> ', where ' <i>a</i> ' > ' <i>b</i> ' |
| P-93.5.3.3 | Specification of double bonds |

| | |
|-------------|---|
| P-93.5.3.4 | Specification of nontetrahedral stereogenic centers |
| P-93.5.3.5 | Axial chirality of spiro compounds |
| P-93.5.4 | Fused and bridged fused compounds |
| P-93.5.4.1 | Specification of configuration by CIP stereodescriptors |
| P-93.5.4.2 | The descriptors ' <i>cisoid</i> ' and ' <i>transoid</i> ' |
| P-93.5.5 | Cyclophanes |
| P-93.5.5.1 | Specification of stereogenic planes |
| P-93.5.5.2 | Specification of chirality centers |
| P-93.5.5.3 | Specification of double bonds |
| P-93.5.6 | Fullerenes |
| P-93.5.6.1 | Definitions and general methodology |
| P-93.5.6.2 | Type 1. Inherently chiral parent fullerenes |
| P-93.5.6.3 | Type 2. Substituted fullerenes inherently chiral because of their substitution pattern |
| P-93.5.6.4 | Type 3. Substituted fullerenes noninherently chiral because of their substitution pattern |
| P-93.5.6.5 | Type 4. Chirality due to chiral substituents |
| P-93.5.6.6 | Superimposition of stereogenic elements in a fullerene molecule |
| P-93.5.7 | Ring assemblies |
| P-93.5.7.1 | Specification of stereogenic axes |
| P-93.5.7.2 | Specification of stereogenic centers |
| P-93.5.7.3 | Specification of double bonds in unsaturated rings and ring systems |
| P-93.6 | Compounds Composed of Rings and Chains |
| P-94 | Conformation and Conformational Stereodescriptors |
| P-94.1 | Definition |
| P-94.2 | Torsion Angle |
| P-94.2.1 | Torsion angle 'X-A-B-Y' |
| P-94.2.2 | If 'A' and 'B' in 'X-A-B-Y' are trigonal centers, a lone pair of electrons |
| P-94.3 | Specific Stereodescriptors |
| P-94.3.1 | Eclipsed, staggered and gauche (or skew) conformations |
| P-94.3.2 | Stereodescriptors for denoting the conformations of alicyclic rings |

P-10 PARENT STRUCTURES FOR NATURAL PRODUCTS AND RELATED COMPOUNDS

| | |
|--------------|--|
| P-100 | Introduction |
| P-101 | Nomenclature for Natural Products Based on Parent Hydrides (Alkaloids, Steroids, Terpenes, Carotenes, Corrinoids, Tetrapyrroles, and Similar Compounds) |
| P-101.1 | Biologically Based Trivial Names |
| P-101.1.1 | Trivial name of a compound isolated from a natural source |
| P-101.1.2 | The ending 'une' or 'iune' |
| P-101.2 | Semisystematic Nomenclature for Natural Products (Stereoparent Hydrides) |
| P-101.2.0 | Introduction |

| | |
|-----------|--|
| P-101.2.1 | General guidelines for choosing a parent structure |
| P-101.2.2 | Structural features allowed for parent structures |
| P-101.2.3 | Numbering of parent structures |
| P-101.2.4 | Identification of individual rings |
| P-101.2.5 | Atomic connector, terminal segment and bond connector |
| P-101.2.6 | Stereochemical configuration of parent structures |
| P-101.2.7 | Semisystematic names of recommended fundamental parent structures |
| P-101.3 | Skeletal Modifications of Parent Structures |
| P-101.3.0 | Introduction |
| P-101.3.1 | Removal of skeletal atoms without affecting the number of rings |
| P-101.3.2 | Addition of skeletal atoms without affecting the number of rings |
| P-101.3.3 | Bond formation |
| P-101.3.4 | Bond cleavage |
| P-101.3.5 | Bond migration |
| P-101.3.6 | Removal of a terminal ring |
| P-101.3.7 | Combination of the prefixes 'cyclo', 'seco', 'apo', 'homo', and 'nor' |
| P-101.4 | Replacement of Skeletal Atoms |
| P-101.4.1 | General methodology |
| P-101.4.2 | Skeletal replacement of carbon atoms by heteroatoms |
| P-101.4.3 | Skeletal replacement of heteroatoms by carbon atoms |
| P-101.4.4 | Skeletal replacement of heteroatoms by other heteroatoms |
| P-101.4.5 | Indicated hydrogen |
| P-101.5 | Addition of Rings and Ring Systems |
| P-101.5.1 | Mannich rings and ring systems incorporated by fusion nomenclature |
| P-101.5.2 | Rings and ring systems incorporated by bridged fused ring nomenclature |
| P-101.5.3 | Rings and ring systems incorporated by spiro nomenclature |
| P-101.6 | Modification of the Degree of Hydrogenation of Parent Structures |
| P-101.6.1 | By changing 'an' or 'ane' to 'ene' or 'yne' |
| P-101.6.2 | The descriptors ' <i>E</i> ' and ' <i>Z</i> ' |
| P-101.6.3 | The prefix ' <i>all</i> ' is used in front of stereodescriptors |
| P-101.6.4 | Double bonds in a parent structure modified by the prefix 'hydro' |
| P-101.6.5 | Carbocyclic or heterocyclic ring components fused to a parent structure saturated using 'hydro' prefixes |
| P-101.6.6 | Additional unsaturation in a parent structure by the prefix 'dehydro' |
| P-101.6.7 | Rearrangement of double bonds by a combination of 'hydro' and 'dehydro' prefixes |
| P-101.7 | Derivatives of Parent Structures |
| P-101.7.1 | The suffixes and prefixes of organic nomenclature and the stereodescriptors α , β , and ξ |
| P-101.7.2 | Modifications of principal characteristic groups |
| P-101.7.3 | Substituent groups derived from parent structures by the suffixes 'yl', 'ylidene', or 'ylidyne' |
| P-101.7.4 | Addition of rings denoting functional groups |

| | |
|--------------|--|
| P-101.7.5 | The prefix 'de' (not 'des'), followed by the name of a group or atom |
| P-101.8 | Further Aspects of Configurational Specification |
| P-101.8.1 | Inversion of configuration |
| P-101.8.2 | Racemates |
| P-101.8.3 | Relative configuration |
| P-101.8.4 | The stereodescriptors 'R' and 'S' for absolute configuration of an achiral parent structure and after ring opening |
| P-102 | Carbohydrate Nomenclature |
| P-102.1 | Definitions |
| P-102.1.1 | Carbohydrates |
| P-102.1.2 | Monosaccharides |
| P-102.1.3 | Oligosaccharides |
| P-102.1.4 | Polysaccharides |
| P-102.2 | Parent Monosaccharides |
| P-102.2.1 | Parent aldoses and ketoses with up to six carbon atoms |
| P-102.2.2 | Numbering parent structures |
| P-102.3 | Configurational Symbolism |
| P-102.3.1 | The Fischer projection of the acyclic form |
| P-102.3.2 | The stereodescriptors 'D' and 'L' |
| P-102.3.3 | The configurational atom |
| P-102.3.4 | Cyclic forms of monosaccharides |
| P-102.3.5 | Conformation of monosaccharides |
| P-102.3.6 | The Mills depiction |
| P-102.3.7 | Stereodescriptors for denoting racemates and uncertain configurations |
| P-102.4 | Choice of a Parent Structure |
| P-102.5 | Monosaccharides: Aldoses and Ketoses; Deoxy and Amino Sugars |
| P-102.5.1 | Aldoses |
| P-102.5.2 | Ketoses |
| P-102.5.3 | Deoxy sugars |
| P-102.5.4 | Amino sugars |
| P-102.5.5 | Thio sugars |
| P-102.5.6 | Substituted monosaccharides |
| P-102.5.6.1 | O-Substitution |
| P-102.5.6.2 | Glycosides |
| P-102.5.6.3 | C-Substitution |
| P-102.5.6.4 | N-Substitution |
| P-102.5.6.5 | Alditols |
| P-102.5.6.6 | Monosaccharide carboxylic acids |
| P-102.5.6.7 | Anhydrides |
| P-102.6 | Monosaccharides and Derivatives as Substituent Groups |
| P-102.6.1 | Glycosyl groups |
| P-102.6.2 | Substituent groups other than glycosyl groups |

| | |
|--------------|---|
| P-102.7 | Disaccharides and Oligosaccharides |
| P-102.7.1 | Disaccharides |
| P-102.7.2 | Oligosaccharides |
| P-103 | Amino Acids and Peptides |
| P-103.0 | Introduction |
| P-103.1 | Names, Numbering and Configuration Specification of Amino Acids |
| P-103.1.1 | Retained and systematic names |
| P-103.1.1.1 | Retained names of the 'common' amino acids |
| P-103.1.1.2 | Retained names of 'less common' amino acids |
| P-103.1.1.3 | Systematic substitutive names |
| P-103.1.2 | Numbering of α -amino carboxylic acids |
| P-103.1.3 | Configuration of α -amino carboxylic acids |
| P-103.2 | Derivatives of Amino Acids |
| P-103.2.1 | System for denoting locants |
| P-103.2.2 | Names of substituent groups |
| P-103.2.3 | Derivatives formed by substitution |
| P-103.2.4 | Ionization of characteristic groups |
| P-103.2.5 | Acyl groups |
| P-103.2.6 | Esters |
| P-103.2.7 | Amides and other nitrogenous derivatives |
| P-103.2.8 | Alcohol, aldehydes, ketones and nitriles |
| P-103.3 | Nomenclature of Peptides |
| P-103.3.1 | Definitions |
| P-103.3.2 | Names of peptides |
| P-103.3.3 | Symbols of peptides |
| P-103.3.4 | Indication of configuration in peptides |
| P-103.3.5 | Modification of named peptides |
| P-103.3.5.1 | Replacement of residues |
| P-103.3.5.2 | Extension of the peptide chain |
| P-103.3.5.3 | Insertion of residues |
| P-103.3.5.4 | Removal of residues |
| P-103.3.6 | Cyclic peptides |
| P-104 | Cyclitols |
| P-104.0 | Introduction |
| P-104.1 | Definitions |
| P-104.2 | Name Construction |
| P-104.2.1 | Stereoisomeric inositols |
| P-104.2.2 | Cyclitols other than inositols |
| P-104.2.3 | Numbering on cyclitols |
| P-104.3 | Derivatives of Cyclitols |
| P-104.3.1 | Derivatives of inositols |
| P-104.3.2 | Derivatives of cyclitols other than inositols |

| | |
|--------------|--|
| P-105 | Nucleosides |
| P-105.0 | Introduction |
| P-105.1 | Retained Names of Nucleosides |
| P-105.2 | Substitution on Nucleosides |
| P-105.2.1 | Substitution of nucleosides |
| P-105.2.2 | Nomenclature when a higher priority characteristic group is present |
| P-106 | Nucleotides |
| P-106.0 | Introduction |
| P-106.1 | Retained Names |
| P-106.2 | Nucleoside Diphosphates and Triphosphates |
| P-106.3 | Derivatives of Nucleotides |
| P-106.3.1 | Nucleotides having retained names |
| P-106.3.2 | Analogues of nucleoside di- and polyphosphates |
| P-106.3.3 | Nomenclature when a higher priority characteristic group is present |
| P-106.3.4 | Oligonucleotides |
| P-106.3.5 | Nucleotides with phosphorothioic acid |
| P-107 | Lipids |
| P-107.0 | Introduction |
| P-107.1 | Definitions |
| P-107.2 | Glycerides |
| P-107.3 | Phosphatidic Acids |
| P-107.3.1 | Phosphatidic acids |
| P-107.3.2 | Configuration of phosphatidic acids |
| P-107.3.3 | Phosphatidylserines |
| P-107.3.4 | Phosphatidylcholines |
| P-107.3.5 | Phosphatidylethanolamine |
| P-107.3.6 | Phosphatidylinositols |
| P-107.4 | Glycolipids |
| P-107.4.1 | Definitions |
| P-107.4.2 | Glycoglycerolipids |
| P-107.4.3 | Glycosphingolipids |
| References | |
| Appendix 1 | Seniority list of elements and ‘a’ terms used in skeletal replacement (‘a’) nomenclature |
| Appendix 2 | Detachable Prefixes |
| Appendix 3 | Structures for Alkaloids, Steroids, Terpenoids, etc. |
